

Biology and Medicine

Open Access





Biology and Medicine Journal being a broad discipline in the Medical research field offers the best platform to publish innovative ideas and researches. It is an indexed bimonthly journal that aims to showcase outstanding research articles from all areas of biology and medicine.



Biology and Medicine Journal is an International, Open- Access, Peer-reviewed, comprehensive journal, publishing the most interesting and advanced research findings in the scientific community. The motto of the journal is to publish original, genuine and relevant articles in the current research field.



The focus broadly covers Cellular biology, Developmental biology, Medical sciences, human biology, Medical biology, Anatomy, Physiology, Disease research etc.

Biology and Medicine - Open Access uses online manuscript submission, review and tracking systems for quality and quick review processing. Submit your manuscript at

www.editorialmanager.com/medsciences

Editors & Editorial Board



Amina S Woods
National Institute of Health, USA



Simon Kaja
University of Missouri USA



Jimmy Thomas Efrid
University of North Carolina, USA



Jamal Zweit
Virginia Commonwealth University USA



Jagat R Kanwar
Deakin University Australia



Tao Zhang
The Research Institute for Children United States



Yujing Li
Emory University School of Medicine USA



Mohammad Afzal
Professor Zoology Aligarh



Anita Mukherjee
Professor University of Calcutta, India



Y Hasan Siddique
Aligarh Muslim University, India



C Perry Chou
University of Waterloo Canada



Kaiser Jamil
Centre for Biotechnology and Bioinformatics, India



N Pradhan
National Institute of Mental Health and Neurosciences, India



Ninoslav Djelic
University of Belgrade Serbia



Qamar Rahman
Integral University India



Ramesh K Aggarwal
Deputy Director, Centre for Cellular and Molecular Biology Hyderabad India



Alan Prem kumar
University of North Texas, USAff



Sangeeta Shukla
Sangeeta Shukla India



Suhel Parvez
Associate Professor India



Tapasya Srivastava
University of Delhi South Campus India



VV Pillay
Amrita Institute of Medical Sciences & Research, India



Anupam Bishayee
American University of Health Sciences, USA



Alejandro Giorgetti
University of Verona Italy



Idress Hamad A
Omar El-Mukhtar University, Libya



Zewei Miao
Agricultural Engineering Sciences Building Urbana



HS Abou-Auda
King Saud University Saudi Arabia



M Jahangir Alam
University of Houston College of Pharmacy USA



M Salah Khalil
Central of Agricultural Pesticides Laboratory Egypt



Haroon Khan
Gandhara University Pakistan



Jibing Yang
University of Michigan USA



Amin A Fadl
University of Wisconsin-Madison, USA



Wei Li
Novartis Institutes for Biomedical Research China



Pierluigi Scalia
Azusa Pacific University, California



Jukka Hartikka
San Diego, USA



David R Bachinsky
Molecular Creativity USA



Medani P Bhandari
Syracuse University New York

OMICS International

5716 Corsa Ave, Suite 110, Westlake, Los Angeles, CA 91362-7354, USA, E-mail: editor.blm@omicsonline.net
Phone: +1-650-268-9744, Fax: +1-650-618-1414, Toll free: +1-800-216-6499

The Role of Telomeres in Cancer Development and Progression, and the Double Edge Sword Effect with Tamoxifen

Nezami MA^{1*}, Hager S² and Garner J¹

¹Research Cancer Institute of America, Pacific Medical Center of Hope, Fresno, CA 93720, USA

²Ccare, Fresno, CA, USA

Keywords: Tamoxifen; Cancer prevention; Telomeres

Introduction

The current practice regarding the use of estrogen receptor antagonists in breast cancer, (such as Tamoxifen), has opened doors for investigators to research these drugs for their use in the prevention of breast cancer. Although this approach is suggested by the literature to be an effective strategy, currently there are no established guidelines applicable to the use of anti-estrogen receptor drugs or prevention in the clinical setting [1,2].

We review the literature related to Tamoxifen therapy and recommend expansion of the application of low dose Tamoxifen for patients with positive estrogen receptor cancers to preventive oncology. These applications include patients at risk for hormonally responsive cancers, and expand beyond that to individuals who may not be even at risk as defined by current risk stratification standards, such as strong family history of breast cancer. In this article we also review an opposing negative effect of Tamoxifen in patients with active cancer, and discuss why we believe it should be “avoided” in the treatment of solid tumors (except breast cancer), although it may be a meaningful tool in prevention of cancer of all types.

Background

The double edge sword of Tamoxifen (favorable in prevention and unfavorable in therapy), is based on our novel understanding of telomeres and their role in carcinogenesis and cancer progression [3-6]. Further studies are warranted in application of low dose Tamoxifen to prevent all solid tumors in unaffected individuals, in a safe and efficacious manner. The theory of how Tamoxifen interacts with telomeres also provides a basic understanding of possible unfavorable mechanisms involved with Tamoxifen’s application and its resistance in treated tumors.

Telomeres and Tamoxifen

Tamoxifen is shown in the literature to be a telomerase activator [7]. The activation of telomeres is an important step in carcinogenesis as it is secondary to what is known as a “telomere crisis” [3,8]. As we age, the length of the telomeres shortens. Continuation of this process over time is important, as it causes the cell DNA to be exposed to genomic instability. This instability at a critical length of telomeres promotes apoptosis, by activating the P53 and other onco-suppressor genes in normal circumstances.

However, if this instability continues beyond a certain point, and/or if the P53 system is impaired, or if the cell is exposed to a telomerase activator (such as Tamoxifen), it can cause DNA damage and mutations in both oncogene promoters as well as oncogene suppressors, which can promote the cancer development at the first stages (carcinogenesis). It is therefore postulated that the telomere crisis happens long before the DNA mutations required for carcinogenesis. The unstable DNA shares the same alterations identified in cancer. Finally, activation of telomerase can save the cell from apoptosis, and death. This activation is uniform in almost all

cancer cells, with high proliferative index [4] shown by both tumor tissue biopsy as well as liquid biopsies. Although it is suggested that telomerase activity is not required for malignant transformation, and it merely an index for highly proliferative cells, it is commonly used as a marker for detection and monitoring the cancer cells, both in tissue biopsies as well as circulating tumor cells or CTCs (Telomerase is an indicator for presence of tumor epithelial cells in the blood) [9,10].

Studies have shown that by using tamoxifen in a cellular culture, the ability of the cells to activate telomerase increases, and as the result the cell will continue to stay young, with prolonged resistance to crisis [7]. This can prevent the cell from an increased amount of genomic instability and further carcinogenesis. However, the same mechanism can explain the cancerous cell’s ability to survive longer in the presence of Tamoxifen, as a telomerase activator [7]. The cancer cells exposed to 4-hydroxy Tamoxifen in culture are able to survive longer and infiltrate distant organs in mice studies, explaining the increased invasion. The critical question remains whether the application of Tamoxifen as early as possible in individuals not known to have cancer may prevent the telomere crisis, and if so, it can prevent carcinogenesis long before it is clinically diagnosed.

Tamoxifen therapy also may have unintended effects on gene expression. This theory is presented in the following two cases studies, both patients with breast cancer who had positive circulating tumor DNA for Neurofibromatosis-I gene post tamoxifen therapy.

Case Studies

Breast cancer with positive NF-1

70 year old female with history of left breast ductal carcinoma diagnosed in 2001 status post lumpectomy and hormonal therapy. Hormonal therapy consisted of Femara for 3 years and Arimidex for 4 years. She was also treated with radiation, and was status-post recurrence of the tumor in same side documented in September 2013, through core biopsy and referral for surgery (mastectomy), and chemotherapies. The extent of disease was documented to involve the skin and required neoadjuvant chemotherapies. Her left breast had a 5 cm mass. She was seen and evaluated for complimentary medicine, and immediately started on IV epigenetic therapies after initial labs.

***Corresponding author:** Nezami MA, Research Cancer Institute of America, Pacific Medical Center of Hope, Fresno, CA 93720, USA, Tel: (559) 475 4300; E-mail: amnezami@yahoo.com

Received: February 23, 2016; **Accepted** March 18, 2016; **Published** March 21, 2016

Citation: Nezami MA, Hager S, Garner J (2016) The Role of Telomeres in Cancer Development and Progression, and the Double Edge Sword Effect with Tamoxifen. Biol Med (Aligarh) 8: 296. doi: [10.4172/0974-8369.1000296](https://doi.org/10.4172/0974-8369.1000296)

Copyright: © 2016 Nezami MA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Her initial labs showed extensively elevated angiogenic markers, including Interleukin 8 (both in serum and plasma). Her Interleukin 8 was 103.3 in plasma (normal less than 34) and 136 in serum (normal less than 66), measured on 6/3/14. Treatment was given every day in first two weeks, five days a week, and twice a week after. She was started on Tamoxifen at a dose of 5 mg daily. Her labs were repeated on 6/17/14 and 8/29/14. She reported improved QOL while receiving therapies and no side effects experienced. Her labs showed marked reduction in both markers. Her IL-8 in serum dropped down to 89 and in her plasma down to 16.8. Further this dropped down to normal range. Her Interleukin 8 normalized both in serum as well as plasma (50 in serum and 27 in plasma), on 10/8/2014.

She subsequently underwent surgery on 12/14, and currently is in remission. She received Firmagon injection, as well as Cytosin (metronomic dose) along with our therapy, and her VEGF dropped down to less than 9, on 5/28/15. Her circulatory tumor DNA (ctDNA) was studied through Guardant 360 Lab, on 6/25/15, which showed positive NF-1, and APC, likely due to the exposure to anti-estrogen therapy. Following further treatments with MTET (Multi-targeted Epigenetic therapy), her Guardant 360 was repeated and showed no ctDNA (Figure 1).

This anti-angiogenetic response to the therapies in a novel approach



Figure 1: Report showing the case study of case 1.

needs to be investigated further in clinical trials to reduce the risk of recurrence and improve response to conventional treatments.

Breast cancer with positive NF-1

43 year old female with history of breast ductal carcinoma, 100 percent ER and PR positive, and a Her-2 negative. Her Ki67 was 80 percent; SBR on biopsy was determined to be 8/9.

She had exhausted many alternative cancer therapies since 5/2012,

when she was diagnosed first. None of the therapies was effective, and in fact her cancer grew to 4 cm, and she opted for bilateral mastectomy and axial lymph node resection in November 2012, which was positive for 2 LN invasions. Upon her arrival at our clinic, she had recurrence of her tumor, with two large palpable sites in the chest on skin. She was complaining of a new onset consistent and dry cough, and was concerning for lung metastasis. Upon evaluation, we ordered a PET scan, which showed multiple pulmonary metastasis bilaterally, which were too numerous to count, with an SUV max of 4.8. She had multiple



Figure 2: Report showing the case study of case 2.

masses in right breast with max SUV of 12.9, and multiple smaller lesions with max SUV of 5.5, plus mediastinal and internal mammary LNs in chest and supraclavicular LNs. In addition she had multiple metastatic LNs in right axilla with SUV max of 2.9.

She had tumor markers all elevated in labs. Immediately she was started on IV epigenetic therapies which she received on daily basis. After 10 treatments, her labs were rechecked. The results showed significant decreases in all her tumor markers. Her cough improved, and she maintained her treatments at our clinic. She did not change her diet, nor did she start any chemotherapy. At this time, she was not on any hormonal blockade, confirming the independent response to epigenetic therapy. In November 2014, she had a restaging PET scan, showing marked reduction of the activities in mediastinal and internal mammary LNs, with improved diffuse pulmonary metastasis. The PET scan was performed and compared with PRE treatment scan after 6 weeks of therapy. The following PET scan done on 1/12/15, demonstrated complete resolution of all pulmonary metastasis and axillary/supraclavicular lymph nodes. There was minimal residual disease reported in the mediastinal lymph nodes, with SUV activity decreased to 5 from 10.6, and to 3 from 7. At this time, the tumor markers were substantially reduced and most returned to normal limits. CA 15.3 was at 31, previously 37, 52, and 56 (respective dates: 4/7/15, 1/16/15, 11/21/14, 10/19/14). CA 27.29 was 33, previously 44, 47, 55.6, and 65 (respective dates 11/21/14, 10/9/14, 9/11/14, 8/8/14). CA 125 was 18.6, previously 19.5 and 39.8 (respective dates 4/7/15, 1/16/15 and 10/9/14). At this time she was receiving combination epigenetic therapies with hormonal blockade, consisting of GnRH blockade (Lupron) as well as Tamoxifen at 5 mg every other day (1/8 normal dose of 20 mg per day). She was restaged again with a PET/CT on 4/15/2015, which showed progression of the local disease in surgical incision, but only minimal residual disease and faint FDG activity (SUV activity of 1.5 compared to 4.8 pre-treatment) in her mediastinal pulmonary lesions, with complete resolution of liver findings. Due to the fact that she was down staged from a Stage IV disease to a local disease in the breast, it was justified to remove the breast tissue and treat her locally with surgery.

Following her surgery, her circulatory tumor DNA was evaluated through Guardant 360, which revealed positive EGFR. Subsequently the patient was started on Tarceva, and the Guardant 360 test was repeated. Although the EGFR mutation allele fraction (MAF) decreased with therapy, the second test showed NF-1, ALK, EGFR and cMET (Figure 2). The presence of NF-1 confirmed the mutated oncosuppressor gene, likely due to the exposure to anti-estrogen therapy, an example of induced heterogeneity. The presence of other alterations, confirmed the induction of heterogeneity by Tarceva. The presence of NF-1 and other alterations, although considered poor prognostic markers, did not change this patient's outcome, or her disease free survival. We believe that this success was due to the combination of epigenetic therapies.

This case suggests an independent, as well as synergistic method of epigenetic therapies combined with hormonal blockade, which was able to treat non-operable Stage IV breast cancer, and down stage the patient successfully to treat the local disease. This quick response

suggests further need to evaluate this novel combinational approach in treating resistant and recurrent advanced breast cancer.

Conclusion

Tamoxifen has been proven to prevent invasive and noninvasive breast cancer [1,2]. Retrospective studies also indicate an effect on incidence of endometrial and ovarian cancer [11,12]. We also recommend large scale studies using Tamoxifen for prevention of cancer. Perhaps the most important question that remains to be answered is the timing of Tamoxifen administration in the prevention of cancer.

References

1. Powles TJ, Ashley S, Tidy A, Smith IE, Dowsett M (2007) Twenty-Year Follow-up of the Royal Marsden Randomized, Double-Blinded Tamoxifen Breast Cancer Prevention Trial. *J Natl Cancer Inst* 99: 283-290.
2. Cuzick J, Forbes JF, Sestak I, Cawthorn S, Hamed H, et al. (2007) Long-term results of tamoxifen prophylaxis for breast cancer--96-month follow-up of the randomized IBIS-I trial. *J Natl Cancer Inst* 99: 272-282.
3. Blackburn EH (2005) Telomerase and Cancer: Kirk A. Landon--AACR prize for basic cancer research lecture. *Mol Cancer Res* 3: 477-482.
4. Belair CD, Yeager TR, Lopez PM, Reznikoff CA (1997) Telomerase activity: a biomarker of cell proliferation, not malignant transformation. *Proc Natl Acad Sci USA* 94: 13677-13682.
5. Lu L, Zhang C, Zhu G, Irwin M, Risch H, et al. (2011) Telomerase expression and telomere length in breast cancer and their associations with adjuvant treatment and disease outcome. *Breast Cancer Res* 13: R56.
6. Wang Z, Kyo S, Maida Y, Takakura M, Tanaka M, et al. (2002) Tamoxifen regulates human telomerase reverse transcriptase (hTERT) gene expression differently in breast and endometrial cancer cells. *Oncogene* 21: 3517-3524.
7. Henson JD, Neumann AA, Yeager TR, Reddel RR (2002) Alternative lengthening of telomeres in mammalian cells. *Oncogene* 21: 598-610.
8. Blocking Telomerase Kills Cancer Cells but Provokes Resistance, *Progression* (2012) *Drug Week*.
9. Xu T, Lu B, Tai Y, Goldkorn A (2010) A Cancer Detection Platform Which Measures Telomerase Activity from Live Circulating Tumor Cells Captured on a Microfilter. *Cancer Research* 70: 6420-6426.
10. Fizazi K, Morat L, Chauveinc L, Prapotnich D, De Crevoisier R, et al. (2007) High detection rate of circulating tumor cells in blood of patients with prostate cancer using telomerase activity. *Ann Oncol* 18: 518-521.
11. Vrscaj MU, Kovacic J, Bebar S, Djuricic A, Fras P, et al. (2001) Endometrial and other primary cancers after tamoxifen treatment of breast cancer — results of retrospective cohort study. *Eur J Obstet Gynecol Reprod Biol* 95: 105-110.
12. Fisher B, Costantino JP, Wickerham DL, Redmond CK, Kavanah M, et al. (1998) Tamoxifen for Prevention of Breast Cancer: Report of the National Surgical Adjuvant Breast and Bowel Project P-1 Study. *J Natl Cancer Inst* 90: 1371-1388.

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submit/

Citation: Nezami MA, Hager S, Garner J (2016) The Role of Telomeres in Cancer Development and Progression, and the Double Edge Sword Effect with Tamoxifen. *Biol Med (Aligarh)* 8: 296. doi: [10.4172/0974-8369.1000296](https://doi.org/10.4172/0974-8369.1000296)